

## Cold ion chemistry within a Rydberg electron orbit: the effect of the molecular structure at the lowest collision energies

V. G. Zhelyazkova<sup>1</sup>, F. B. V. Martins<sup>1</sup>, S. Schilling<sup>1</sup>, J. A. Agner<sup>1</sup>, H. Schmutz<sup>1</sup>, F. Merkt<sup>1\*</sup>

<sup>1</sup>ETH Zurich, Department of Chemistry and Applied Biosciences, Vladimir-Prelog-Weg 2, 8093 Zürich, Switzerland

In the study of ion-molecule reactions, Rydberg atoms and molecules can be used as a proxy for the ions. The Rydberg electron shields the ion-core-molecule reaction from stray electric fields, making it possible to reach collision energies ( $E_{\text{coll}}$ ) near zero Kelvin [1,2]. Rydberg atoms can possess high induced electric dipole moments which allows for the manipulation of their translational motion by modest electric-field gradients using surface-electrode devices [3]. We use such a device to merge a beam of Rydberg helium atoms [He( $n$ )] with a molecular supersonic beam. We vary the relative velocity between the reactants by changing the velocity of the He( $n$ ) atoms, and thereby the collision energy. With this method, we can reach  $E_{\text{coll}}/k_{\text{B}}$  from  $\sim 40$  K all the way down to  $\sim 100$  mK in the moving centre-of-mass reference frame.

In this contribution, we present an overview of the results of experimental studies of reactions between the He<sup>+</sup> ion and several small molecules. We find that for molecules with a strong electric dipole moment (such as CH<sub>3</sub>F [4] and NH<sub>3</sub> [5]) the reaction yield is strongly enhanced near 0 K, and the reaction rate coefficients reach more than ten times the typically assumed Langevin rate coefficient. For molecules with no dipole moment (e.g. N<sub>2</sub> [6]), the opposite effect is observed, *i.e.*, a *suppression* of the reaction yield near 0 K. We also investigate the reactions with molecules which have only a small dipole moment and find that the  $E_{\text{coll}}$ -dependence of the reaction yields in this case are determined by the character of the electronic ground state. For example, we measure strikingly different reaction yields for NO (<sup>2</sup> $\Pi$  character [7]) and CO (<sup>1</sup> $\Sigma^+$  character [8]). Our experimental results are interpreted with the aid of a rotationally-adiabatic-capture model [5,6] inspired by earlier theoretical treatments [9,10].

[1] M. Matsuzawa, *Phys. Rev. A*, **2010**, 82, 054701.

[2] F. B. V. Martins, V. Zhelyazkova, Ch. Seiler, F. Merkt, *New J. Phys.*, **2021**, 23, 095011.

[3] P. Allmendinger, J. Deiglmayr, J., J. A. Agner, H. Schmutz, H. and F. Merkt, *Phys. Rev. A*, **2014**, 90, 043403.

[4] V. Zhelyazkova, F. B. V. Martins, J. A. Agner, H. Schmutz, F. Merkt, *Phys. Rev. Lett.*, **2020**, 125, 26340.

[5] V. Zhelyazkova, F. B. V. Martins, J. A. Agner, H. Schmutz, F. Merkt, *Phys. Chem. Chem. Phys.*, **2021**, 23, 21606.

[6] V. Zhelyazkova, F. B. V. Martins, M. Žeško, F. Merkt, *Phys. Chem. Chem. Phys.*, **2022**, 24, 2843.

[7] V. Zhelyazkova, F. B. V. Martins, S. Schilling, F. Merkt, *J. Phys. Chem. A*, **2023**, 127, 1458.

[8] F. B. V. Martins, V. Zhelyazkova, F. Merkt, *New J. Phys.*, **2022**, 24, 113003.

[9] J. Troe, *J. Chem. Phys.*, **1987**, 87, 2773.

[10] D. C. Clary, *Ann. Rev. Phys. Chem.*, **1990**, 41, 61.