

## Encoding Stereochemistry in Molecular Fingerprints

M. Orsi<sup>1</sup>, D. Probst<sup>2</sup>, J. Reymond<sup>3\*</sup>

<sup>1</sup>University of Bern, Department of Chemistry, Biochemistry and Pharmacy, <sup>2</sup>École Polytechnique Federale de Lausanne, <sup>3</sup>University of Bern

Molecular fingerprints are important tools enabling fast similarity comparisons between molecules<sup>1</sup>. These fingerprints encode different molecular properties, such as pharmacophores<sup>2</sup>, atom distances<sup>3,4</sup> and substructures<sup>4-6</sup>, and are particularly useful in the context of virtual screening, where one carries out millions of comparisons to detect potential hits. However, a serious limitation of the most well-known fingerprints is their inability to account for chirality in a chemically meaningful manner. Specifically, the challenge lies in encoding chirality to distinguish enantiomers and diastereomers as highly similar yet distinct entities. In this study, we introduce MHFP6\* and MAP4\*, enhanced versions of our MHFP6<sup>6</sup> and MAP4<sup>4</sup> fingerprints, which correctly encode CIP chirality in a chemically meaningful manner. Both fingerprints demonstrate the capability to correctly differentiate isomers, thereby overcoming the previously mentioned limitations.



[1] Peter Willet, *Drug Discov. Today* **2006**, *11* (23-24), 1046-1053.

[2] Alice Capecchi, Mahendra Awale, Daniel Probst, Jean-Louis Reymond, *Mol. Inform.* **2019**, *38* (5), 1900016.

[3] Raymond E. Carhart, Dennis H. Smith, Rengachari Venkataraghavan, *J. Chem. Inf. Comput. Sci.* **1985**, *25* (2), 64-73.

[4] Alice Capecchi, Daniel Probst, Jean-Louis Reymond, *J. Cheminformatics* **2020**, *12* (1), 43.

[5] David Rogers, Mathew Hahn, *J. Chem. Inf. Model.* **2010**, *50* (5), 742-754.

[6] Daniel Probst, Jean-Louis Reymond, *J. Cheminformatics* **2018**, *10* (1), 66.