


Exploring Regio-Selective Spin Interactions: Positional Isomerism and its Influence on Spin Communication in Light-Induced Multi-Spin Systems

M. El Bitar Nehme¹, P. Thielert², M. Mayländer², T. Quintes², M. Franz², S. Zimmermann³, P. Gilch³, M. Rickhaus^{1*}, S. Richert^{2*}

¹Universität Zürich, Department of Chemistry, Winterthurerstraße 190, 8057 Zürich, Switzerland, ²University of Freiburg, Institute of Physical Chemistry, Albertstraße 21, 79104 Freiburg, Germany, ³Heinrich Heine University Düsseldorf, Institute of Physical Chemistry II, Universitätsstraße 1, 40225 Düsseldorf, Germany

Highly versatile photogenerated multi-spin systems are promising candidates to explore the factors governing spin communication on a molecular level.¹ While the radical acts as a sensitizer that improves the intersystem crossing rate, the delicate covalent linkage between chromophore-radical systems serves as a means of controlling the excited state dynamics of the chromophore.² The aim of this project is to develop covalent multi-spin systems to study spin-information transfer and storage. This is performed by engineering systems that consist of at least two organic spin centers that we connect by a conjugated framework. The bridge between the two spin centers is then systematically modified to trace the changes in the resulting spin communication. By choosing a bridged biphenyl as the linker between the chromophore and the radical, the electronic communication throughout the synthon is expected to vary with the torsion angle Φ between the planes of the two phenyl rings, which in turn modulates the spin-spin interaction.



(1) Yamauchi, S. *Chem. Soc. Jpn.* **2004**, 77 (7), 1255-1268.

(2) Nolden, O.; Fleck, N.; Lorenzo, E. R.; Wasielewski, M. R.; Schiemann, O.; Gilch, P.; Richert, S. *Chem-Eur. J.* **2021**, 27 (8), 2683-2691.