


Efficient alcohol oxidation with novel manganese(III) complexes with bis(phenolate)-NHC ligands

G. Rigoni¹, P. Nylund¹, M. Albrecht^{1*}

¹Department of Chemistry, Biochemistry and Pharmaceutical Sciences, University of Bern, Freiestrasse 3, CH-3012 Bern, Switzerland

Oxidation chemistry plays a critical role in the synthesis of pharmaceuticals and value-added products, such as perfumes. A catalytic approach based on Earth-abundant metal complexes and inexpensive oxidants such as peroxides currently represents the most attractive strategy to perform this transformation.^[1] Our group recently prepared a new class of N-heterocyclic carbene ligands bearing phenolate moieties and their corresponding metal complexes, investigating the potential benefit of the electronic flexibility of the carbene fragment in catalytic applications involving redox events.^[2,3] Here, we decorated imidazole- and triazole-derived carbenes with phenolate substituents^[4,5] to access Mn(III) complexes, which efficiently catalyze the oxidation of alcohols in the presence of *t*BuOOH as terminal oxidant. In addition, we will present the versatility of the catalytic system and mechanistic investigations in order to disclose the nature of the active species.



[1] C. Miao, X.-X. Li, Y.-M. Lee, C. Xia, Y. Wang, W. Nam, W. Sun, *Chem. Sci.* **2017**, 8, 7476–7482.

[2] S. Bertini, M. Rahaman, A. Dutta, P. Schollhammer, A. V. Rudnev, F. Gloaguen, P. Broekmann, M. Albrecht, *Green Chem.* **2021**, 23, 3365–3373.

[3] W. Stroek, M. Keilwerth, D. M. Pividori, K. Meyer, M. Albrecht, *J. Am. Chem. Soc.* **2021**, 143, 20157–20165.

[4] E. Borré, G. Dahm, A. Aliprandi, M. Mauro, S. Dagorne, S. Bellemin-Laponnaz, *Organometallics* **2014**, 33, 4374–4384.

[5] M. Baltrun, F. A. Watt, R. Schoch, C. Wölper, A. G. Neuba, S. Hohloch, *Dalton Trans.* **2019**, 48, 14611–14625.