


Mechanistic Investigations of “Ligand-Free” Kumada-Tamao-Corriu Cross-Coupling Reactions

L. Vedani¹, A. M. Borys¹, E. Hevia^{1*}

¹Universität Bern, Departement für Chemie, Biochemie und Pharmazie

The Kumada-Tamao-Corriu (KTC) cross-coupling reaction was first reported in 1972 and is a powerful synthetic tool to form C-C bonds from Grignard reagents and C(sp²)-halides using nickel catalysts.¹ Only a few years later, in 1979, Wenkert reported the use of aryl ethers as electrophilic coupling partners in KTC reactions.² Since then, the reaction has been further developed and efforts have been directed towards mechanistic understanding.³ The majority of this research however involves the use of bespoke electron-donating ligands, such as NHCs and phosphines.

In this work, we will focus on mechanistic investigations of the KTC reaction using simple Ni(0)-olefin or Ni(II) catalysts, without the use of auxiliary ligands. Under these “ligand-free” conditions, the reaction instead relies on the formation of electron-rich anionic nickelates,⁴ several of which have been isolated and characterised by solution and solid-state techniques. Furthermore, evidence that points towards the importance of bimetallic cooperativity in the Ni-catalysed cross-coupling of aryl ethers will be discussed.⁵



- [1] a) K. Tamao, K. Sumitani, M. Kumada. *J. Am. Chem. Soc.*, **1972**, 94, 12, 4374–4376, b) J. P. Masse, R. J. P. Corriu. *J. Chem. Soc., Chem. Commun.*, **1972**, 144a–144a
- [2] E. L. Michelotti, C. S. Swindell, E. Wenkert. *J. Am. Chem. Soc.*, **1979**, 101, 8, 2246–2247
- [3] J. B. Diccianni, T. Diao. *Trends in Chem.*, **2019**, 1, 9, 830–844
- [4] A. M. Borys, E. Hevia. *Angew. Chem. Int. Ed.*, **2021**, 60, 24659–24667
- [5] L. Vedani, A. M. Borys, E. Hevia. *In preparation*