

Salen-derived multi-metallic complexesJ. Pradegan¹, K. M. Fromm^{1*}¹Department of Chemistry, University of Fribourg, Fribourg, Switzerland

Salen is one of the most studied ligands in chemistry, notably for its straightforward synthesis and its ability to coordinate a variety of metal ions.^[1] This Schiff base compound has a N_2O_2 chelate site allowing the coordination of one metal ion. Salen is an interesting ligand in that it can be functionalized providing a panoply of new ligand structures. The main application of salen-metal complexes is in the field of catalysis,^[2] the most well-known being the Jacobsen catalyst for its enantioselective epoxidation of alkenes.^[3]

In our studies, we aim at synthesizing multi-metallic complexes with two or three different metal ions. To achieve this goal, salen can be functionalized with glycol chains creating e.g. a O_3O_3 recognition site for a second metal ion. The combination of different metal ions can provide enhanced chemical and biological properties.^[2] For instance, the synthesized multi-metallic complexes could be used as mixed metal oxide precursors or show potential synergic antibacterial properties.^[4]




Fig. 1 : Representation of salen-derived ligand with N_2O_2 chelate and O_3O_3 recognition sites

[1] Ashish K. Asatkar, Mamta Tripathi, Deepali Asatkar, *IntechOpen*, **2020**, 1, Ch. 7

[2] Ramananda Maity, Bernhard S. Birenheide, Frank Breher, Biprajit Sarkar, *ChemCatChem*, **2021**, 13, 2337-2370

[3] John Hanson, *J. Chem. Educ.*, **2001**, 78, 9, 1266-1268

[4] Alba Finelli, Sarah-Luise Abram, Nelly Hérault, Aurélien Crochet, Katharina M. Fromm, *Cryst. Growth Des.*, **2020**, 20, 4945-4958