

Nuclear Forensic Investigations of High Purity Depleted Uranium Ammunition

M. Hofstetter^{1,2}, S. Rölli², P. Steinegger^{1,3*}

¹ETH Zürich, Departement für Chemie und Angewandte Biowissenschaften, Vladimir-Prelog-Weg 1, 8093 Zürich, ²Spiez Laboratory, Austrasse, 3700 Spiez, ³Paul-Scherrer Institut, Forschungsstrasse 111, 5232 Villigen

Sample attribution and their comparison are essential skills in a nuclear forensic investigation. Over the past two decades, the analysis of trace element impurities by sector-field inductively coupled plasma mass spectrometry (sf-ICP-MS) has become a valuable source of information. Today, lanthanoid impurities are known to be the most valuable source of information for sample attribution and comparison [1].

In this work, lanthanoid patterns, which have been extensively studied in the field of geology [2,3], are used in a nuclear forensic context to ensure reliable identification and comparison of uranium and thorium materials. Known methods lead to lanthanoid contamination and thus, to the modification of the said patterns [4]. The herein presented method addresses these shortcomings and can be used for the reliable quantification of lanthanoids. The obtained sensitivity allows lanthanoid quantification with a factor of 100 to 1000 lower, in comparison to previously established methods. This allows for a characterization of lanthanoid impurities down to the lower ppb (ng/g) to ppt (pg/g) range. Hence, the developed method is applicable to various high purity uranium materials such as nuclear fuel pellets and depleted uranium ammunition and fills an important analytical gap.

References:

- 1) Elizabeth Keegan et al. *Forensic Science International*, **2014**, 240, 111-121.
- 2) Maria Rita Cicconi et al. *Magma Redox Geochemistry*, Wiley, **2021**, 381-398.
- 3) Wilson, M. *Igneous Petrogenesis*, Springer Netherlands, **2007**.
- 4) Judit Krajc6 et al. *Radiochimica Acta*, **2016**, 104, 471-479.