

## Small Ribozymes in Gut Bacteriophages and Their Potential Role in Microbiome-Virome Interactions

K. Kienbeck<sup>1</sup>, L. Malfertheiner<sup>2</sup>, S. Zelger-Paulus<sup>1</sup>, S. Johannsen<sup>1</sup>, C. von Mering<sup>2\*</sup>, R. K. Sigel<sup>1\*</sup>

<sup>1</sup>Department of Chemistry, University of Zurich, Zurich, Switzerland, <sup>2</sup>Department of Molecular Life Sciences, University of Zurich, Zurich, Switzerland

Hepatitis delta virus (HDV)-like ribozymes belong to the class of small self-cleaving RNA enzymes, which catalyze a site-specific internal transesterification while exhibiting diverse sequences and structures [1]. However, the biological functions of these ribozymes, particularly minimal examples, remain largely unknown [2].

We computationally discovered thousands of minimal HDV-like ribozymes in bacteriophage genomes associated with the human gut. *In vitro* validation was performed on selected examples associated with viral tRNAs, coined theta ribozymes. We propose a novel biological function of these ribozymes, suggesting their involvement in tRNA maturation and the lytic-lysogenic switch of specific phages (Figure 1). Investigating the catalysis and regulation of theta ribozymes sheds light on the poorly understood implications of the microbiome-virome interactions of the human gut and their impact on health and disease.




Figure 1: Proposed phage infection cycle involving theta ribozymes. aaRS: aminoacyl-synthetase; int: integrase;  $\theta$ rz: theta ribozyme; pol: polymerase.

[1] L. Sharmeen, M.Y. Kuo, G. Dinter-Gottlieb, J. Taylor, *Journal of virology* **1988**, 62, 2674-2679.

[2] N.J. Riccitelli, E. Delwart, A. Lupták, *Biochemistry* **2014**, 53, 1616-1626.