


Broadband APT (BAPT): a Versatile APT Experiment with Improved J -Compensation and Optimal Suppression of Artifacts in C_q -only Spectra

P. Bigler¹, D. Chakif¹, I. Gjuroski¹, J. Furrer^{1*}

¹University Bern, Department of Chemistry, Biochemistry and Pharmaceutical Sciences

1D ¹³C-NMR DEPT or APT experiments still belong to the most common experiments for assigning ¹³C signals and for the elucidation of molecular structures on a routine level. The APT sequence¹ suffers from two main drawbacks, found long ago: (i) a low tolerance for wide ranges of ¹J_{CH} values, which in the worst-case cancels signals or produces signals with mistaken multiplicity. (ii) The frequent presence of intense artifacts in the C_q-only spectrum, especially if the range of coupling constants of the investigated molecule is large. Improved APT sequences² (Compensated Attached Proton Test, CAPT2 and CAPT3) have been designed by the group of McClung to improve the tolerance with respect to the wide range of ¹J_{CH} values. The CAPT3 sequence indeed leads to excellent tolerance over a wide range of one-bond J coupling constants.



In this report, we introduce a new APT sequence, the Broadband-APT (BAPT) sequence, which further improves the tolerance of the CAPT sequences to a wide range of ¹J_{CH} values and can provide ultra-clean C_q-only spectra, with CH_n artifact levels as low as those obtained using SEMUT, SEMUT-GL and *i*QCD sequences, known to provide the best C_q-only spectra.

[1] S. L. Patt, J. N. Shoolery, *J. Magn. Reson.* **1982**, *46*, 535–539.

[2] A. M. Torres, T. T. Nakashima, R. E. D. McClung, *J. Magn. Reson. A* **1993**, *101*, 285–294.